

Estimating Alfalfa RFV in the Field Using PEAQ

Increasing alfalfa **Relative Feed Value (RFV)** can add value to the farm enterprise through reductions in purchased feed and dry matter intake. Climatic variations of spring conditions, and its varied impact on alfalfa growth and development makes it impossible to use a calendar date to predict when best to harvest first crop alfalfa. However, research from the University of Wisconsin has created a unique and simple method to predict RFV of standing forage, and estimate the best time to take the first harvest of the season.

It is recommended to harvest alfalfa at approximately 150 RFV for milking dairy herds and 120 to 130 RFV for heifers, stocker cattle and lactating beef cattle. Standing first crop alfalfa can drop 3 to 5 points of RFV per day. A one-week delay of first crop harvest could cost 30 RFV points in feed quality.

The PEAQ procedure estimates the RFV of the standing crop. Under the best conditions, 10 to 15% of the forage dry matter will be lost at harvest. This is typically about 10 to 15 RFV units for haylage, and 20 RFV units for hay. Therefore, it is recommended to cut at 165 to 170 RFV to end up with harvested forage at 150 RFV.

How to PEAQ Your Alfalfa Harvest

Predictive Equations for Alfalfa Quality (PEAQ) is a method to predict the forage quality of standing alfalfa. The two equations predict ADF and NDF when the height of the tallest stem is measured and the maturity of the most advanced plant is determined. The equations have been validated in the Midwest and also in other environments from California to New York. It is a reliable indicator to estimate the optimum harvest time for first-crop alfalfa.

Step 1: Choose a representative area in the field.

Step 2: Determine the most mature stem in the area using the table at the right.

Step 3: Measure the most mature stem in the area. Measure it from the soil surface to the tip of the stem (NOT to the tip of the highest leaf blade). Straighten the stem for an accurate measure of its length. The tallest stem may not be the most mature stem.

Step 4: Based on the most mature stem and length of the tallest stem, use the following chart to determine estimated RFV content of the standing forage.

Step 5: Subtract the 10 to 20 RFV units to account for harvest losses during the haylage or hay harvest process.

--- Stage of Most Mature Stem----

Height of Tallest Stem From soil surface to stem tip.	Late Vegetative No buds visible	Bud Stage 1 or more nodes with visible buds. No flowers visible.	Flower Stage 1 or more nodes with open flower(s)
-Inches -	RFV	RFV	RFV
16	237	225	210
17	230	218	204
18	224	212	198
19	217	207	193
20	211	201	188
21	205	196	183
22	200	190	178
23	195	185	174
24	190	181	170
25	185	176	166
26	180	172	162
27	175	168	158
28	171	164	154
29	167	160	151
30	163	156	147
31	159	152	144
32	155	149	140
33	152	145	137
34	148	142	136
35	145	139	131
36	142	136	128
37	138	133	126
38	135	130	123
39	132	127	121
40	129	124	118
41	127	122	115
42	124	119	113

IOWA STATE UNIVERSITY

University Extension

Fact Sheet BL-22, March 2001

Brian Lang, ISU Extension Crop Specialist,
Larry Tranel, ISU Extension Dairy/Forage Specialist,
adapted from the University of Wisconsin.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Many materials can be made available in alternative formats for ADA clients. To file a complaint of discrimination, write USDA, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964.