

How Do We Estimate N Carryover for 2013?

The drought of 2012 has likely increased the carryover of nitrate-nitrogen (nitrate-N) into the 2013 season. In addition to less N being used by last year's crop, the reduced rainfall in 2012 resulted in less nitrate leaving the soil through leaching and de-nitrification. Although it is common for about 50 pounds per acre of nitrate-N to carryover from one season to the next, soil samples pulled in the fall of 2012 indicate we will likely have fields with more than 100 pounds per acre of carryover nitrate-N into the spring of 2013. *We can take advantage of this carryover N by pulling soil samples this spring to estimate carryover N and reduce spring N rates accordingly for the 2013 corn crop. Failure to do so may result in increased nitrate losses into water systems in 2013 and future years.* To estimate nitrate-N carryover:

1. Pull 1-foot soil samples to at least a 2-foot depth (0-1 foot & 1-2 foot) before the spring N is applied. A 3 foot depth is preferable. Pull 15-30 cores per sample on an area of no more than 10 to 20 acres. Mix thoroughly and send a subsample (standard soil sample size) to the lab to test for nitrate. Multiple samples per field should be collected.
2. Take the soil test result (ppm nitrate-N) times 4 to calculate pounds per acre of N.
3. Add up the N in each foot and subtract the "normal" carryover N (40 pounds per acre for 2 foot depth and 50 pounds per acre for 3 foot depth).
4. Subtract the carryover N from your usual N rate.
5. Regardless of lab results, apply no less than 50 pounds per acre if no N has been applied, to account for field variability.

Example for Estimating Carryover Nitrate-N for 2-Foot Sampling Depth

Assume no N has been applied for 2013 season and 150 pounds per acre is the usual N rate. Assume lab results of 10 ppm nitrate-N for 0-1 foot and 15 ppm nitrate-N for 1-2 foot.

$$\begin{aligned} & \mathbf{150 \text{ lb N/A} - [(10 \text{ ppm} \times 4) + (15 \text{ ppm} \times 4) - 40 \text{ lb N/A}] \\ & \mathbf{= \text{recommended rate of } 90 \text{ lb N/A for 2013 corn}} \\ & \mathbf{150 - [(40+60) - 40] = 90 \text{ lb N/A}} \end{aligned}$$

If there is excess spring rainfall, the carryover N available for the 2013 season will be reduced.

Prepared by Jim Fawcett, ISU Extension Field Agronomist, with information supplied by John Sawyer, ISU Extension Agronomist – Soil Fertility
March, 2013

Cooperative Extension Service, Iowa State University of Science and Technology, and the United States Department of Agriculture cooperating. The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Many materials can be made available in alternative formats for ADA clients. To file a complaint of discrimination, write USDA, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964.