

Monitoring hygiene in calf feeding equipment

Authors:

Jennifer Bentley, Iowa State University Extension and Outreach, Dairy Field Specialist

Ezequías Castillo Lopez, University of Nebraska-Lincoln, Adjunct Professor Dairy Nutrition

Kim Clark, University of Nebraska-Lincoln, Dairy Extension Educator

Paul Kononoff, University of Nebraska-Lincoln, Associate Professor of Dairy Nutrition/Dairy Extension Specialist

Hugo Ramirez Ramirez, Iowa State University, Assistant Professor Dairy Nutrition and Management

Amanda Robinson, Iowa State University, Ph.D. student

Aclaración: Este material se presenta como una guía básica, cada estable debería desarrollar sus Procedimientos Operativos Estandarizados.

Introduction



Calves represent the future of any dairy farm because they will become the next generation of lactating cows. At birth, calves have an immature immune system and are unable to fight off infections by themselves, therefore it is imperative that calves receive high quality colostrum shortly after birth. The colostrum contains proteins called immunoglobulins or antibodies that protect the calf against diseases. These antibodies are produced by the mother but do not transfer to the fetus so they must be obtained through colostrum. In order to start and maintain a

healthy life, calves must be cared for in a way that prevents outbreaks and spread of diseases, and promotes growth and development of their digestive system.

Raising calves involves aspects such as housing, feeding and nutrition, vaccination, and handling. In any of these aspects, it is important to highlight that hygiene is of utmost importance because calves may be susceptible to diseases caused by pathogens harbored in housing facilities, transportation trailers, other calves, or in the feeding equipment. Of these, feeding equipment is perhaps one of the most suitable environments for pathogens to grow, particularly bacteria. Milk or milk replacer are very rich sources of nutrients not only for calves but also for bacteria. Moisture and abundance of nutrients are key elements that favor bacterial



proliferation, therefore, proper sanitation ensures the removal of remaining nutrients and moisture from feeding equipment. The latter is easily achieved by allowing feeding equipment to dry thoroughly. For example, it is highly discouraged to stack wet buckets even if they are put upside-down; instead place them in a drying rack individually. Removal of remaining nutrients is more difficult and harder to evaluate.

Protein, sugars and fat contained in milk or milk replacer can linger on surfaces of feeding equipment such as buckets, bottles, nipples, whisks, mixing jar in automatic calf feeding systems and even on the bristles of


brushes intended to clean bottles. The presence of residue on surfaces is easy to recognize when there is water

beading; when water beading occurs it is likely that the surface has been accumulating residue for a relative long time. When bacteria proliferate they form a layer of residue known as biofilm. This biofilm is a matrix of various substances that allows bacteria to adhere very strongly to a surface. This is commonly see when feeding equipment is simply rinsed with water and mild soap or no soap at all. In order to control the development of biofilm, it is necessary to use proper physical and chemical cleaning practices. Thorough cleaning is accomplished by initially rinsing all equipment with lukewarm water to remove as much milk residue as possible followed by vigorous scrubbing with detergent and water at 120 Fahrenheit (49° C)

Physical	Chemical
Water temperature and contact time	Detergents
Scrubbing and rinsing	Disinfectants
Drying	Sanitizers

It is important to note that each one the chemical agents listed in the table has a very specific function.

Detergents are used to break up organic deposits such as fat and protein, rinsing with water alone is not



effective and bacteria can lie beneath the surface created by residual fat and protein. Once a surface is cleaned with a detergent, a **disinfectant** is used to kill microorganism. Using detergents and disinfectants is an effective combination to reduce the bacterial load and prevent formation of biofilm. A **sanitizer** helps reduce the number of microorganism on a surface but is not as effective as a disinfectant, it may be used a way to improve hygiene in the calf kitchen but not as a substitute of good practices.

Clean facilities and housing is also very important to raise healthy calves. It is recommended to sanitize calf hutches or individual pens before a newborn calf is placed in it, also clean and dry bedding should be provided as frequently as needed.

Monitoring cleanliness in the calf kitchen

Even though surfaces may appear clean to the naked eye, there may be bacteria actively growing and thriving on miniscule amounts of nutritious residue caused by improper cleaning. There are two common ways to evaluate cleanliness of surfaces. One is through the use of a protein swab and the other is through a hand-held device that detects the presence of bacteria via a chemical reaction that produces light, properly known as bioluminescence. Both of these methods are widely used in human food processing facilities to monitor effectiveness of their surfaces and equipment sanitation procedures and prevent the spread of pathogenic bacteria. A calf kitchen is also a food processing facility, therefore, both techniques can be easily applied at the farm level.



Protein swabs This is a quick and economical method to detect the presence of protein on surfaces. If protein is detected, it is an indication of improper removal of residue due to poor physical or chemical cleaning practices. This test is based on a chemical reaction that produces a change in color depending on how much protein residue is present. For this particular example, the results may be:



Prior to the test, the swab must be brought to room temperature. The first step is to twist and remove the swab from the tube, then swab a 4 × 4 inch area in a crisscross pattern applying enough pressure to slightly bend the swab shaft. Re-insert the swab into the tube and hold firmly, snap the valve, squeeze bulb to force the liquid reagent down the tube. Shake for 5-10 seconds, keep the tube upright and wait for the reaction to develop. The quickest result can be seen in 1 minute with highly soiled surfaces and in 10 minutes in cleaner surfaces.

Bioluminescence



This method is used to evaluate the presence of bacteria on surfaces. Like any living organism, bacteria obtain energy from a molecule known as ATP (adenosine triphosphate). After cleaning feeding equipment, a swab is rubbed on a surface. If there are live bacteria present on that surface, the swab would pick up traces of ATP. When the swab is inserted in the luminometer, there is a chemical reaction that produces light and the intensity of the light is measured by the device. The reading is shown in RLU which stands for relative light units. The brighter it is, the more bacteria are present on that surface indicating deficient cleaning practices. It is recommended to work closely with a veterinarian or calf care specialist to determine a base line, caution and correction values as these values may vary by farm depending on the type of surface and cleaning protocols in place.

Iowa State University Extension and Outreach programs are available to all without regard to race, color, age, religion, national origin, sexual orientation, gender identity, genetic information, sex, marital status, disability, or status as a U.S. veteran. Inquiries can be directed to Office of Equal Opportunity, 3350 Beardshear Hall, Ames, Iowa 50011, (515) 294-7612.

IOWA STATE UNIVERSITY
Extension and Outreach



UNL does not discriminate on the basis of race, ethnicity, color, national origin, sex (including pregnancy), religion, age, disability, sexual orientation, gender identity, genetic information, veteran status, marital status, and/or political affiliation in its programs, activities and employment. UNL complies with all local, state and federal laws prohibiting discrimination, including Title IX, which prohibits discrimination on the basis of sex. For more information visit go.unl.edu/nondiscrimination.