

## ISU Extension in Johnson County

3109 Old Hwy 218 S.  
Iowa City, IA 52246  
Phone: (319) 337-2145  
[www.extension.iastate.edu/johnson](http://www.extension.iastate.edu/johnson)

## Table of Contents:

- 1 — Potential Harvest and Storage Challenges
- 2 — Potential Harvest and Storage Challenges (continued)
- 3 — Thatch and Fall Removal
- 3 — Iowa Concern Hotline
- 4 — Fall Commercial Pesticide Schedule
- 4 — Tri-State Ag Lender's Seminar
- 5 — Secure Pork Supply
- 5 — New Youth Program Specialist

## Upcoming Events

Mark your Calendars now with these upcoming events.

**See Page 4** - Pesticide Applicator Training & Certification opportunities

## Plan Ahead for Potential Harvest and Storage Challenges This Fall

Article Author: Kristina TeBockhorst, ISU Extension Ag Engineering Field Specialist, SE IA

In southeast and east-central Iowa roughly half of the corn acres were planted by the first week of May or by mid-May, respectively; the other half of the corn acres were mostly planted the last week of May or early June (Southeast Iowa more so in June).

This year especially, it will be beneficial for farmers to scout fields and plan for harvest, drying, and storing for this fall. Monitor crop development to determine ahead of time whether you might have potential problems with immature, frost-damaged grain and wet grain.

### Potential Challenges:

Frost-damaged soybeans will have a slower dry-down in the field and may produce green/yellow soybeans with above-normal shrink from drying. The green color may subside within two weeks of maturity if allowed to dry in the field or after several weeks of aeration.

Corn damaged by a freeze before it has reached physiological maturity will create issues of low test-weight, low quality, and high-moisture. Light corn has a shorter storage life and is more difficult to dry. Corn with a test weight of 52-pounds or frost-damaged corn will have about half the effective storage life compared to normal quality, 56-pound corn.

Additionally, corn that reaches maturity later in the year can still bring issues of high moisture with less in-field drying between maturity and harvest. Corn grain at maturity, or black layer, is at approximately 30-35% moisture. Corn field drying rate decreases with air temperatures: in September, weekly drying is estimated at 4.5%-points-per-week, and in October, November, and December, this is reduced to 2.5%, 1%, and 0.5%, respectively.

It is too early for accurate estimates of first freeze date, but we can observe historical dates to get an idea of general risk. Use the calculator, found at <https://mrcc.illinois.edu/U2U/gdd/>, along with your location and corn hybrid characteristics to estimate date of corn maturity, or black layer, and see how it compares with historical average first freeze dates.

Using this calculator to observe a situation farmers in southeast or east-central Iowa might be in, we can use a location of Johnson County, Iowa, a planting date of June 1, and corn maturity of 111-days. We also need to manually adjust in the calculator our growing degree day requirements for corn planted after May 1, which requires about 6.8 fewer growing degree days per day to reach maturity. In this situation, the calculator estimates black layer occurring on September 29, 20 days before the average first freeze for this location, or October 20. For the same corn hybrid, moving back to a planting date of June 10, however, the calculator predicts black layer on October 21.

The combination of planting date and actual date of the first freeze this fall will determine if you may be in a situation with increased drying requirements, or if you might be dealing with low-quality, immature grain as well. It is also important to remember that even if you mature in plenty of time before the average first freeze date, there is still a chance of getting an earlier-than-average first freeze.

Article continues on page 2...

## Plan Ahead for Potential Harvest and Storage Challenges This Fall (continued)

### Handling and Storage Recommendations:

Moisture meters are typically inaccurate at high grain moisture levels, so take readings to be rough estimates. When testing for moisture content, be sure to follow the manufacturer's procedure for obtaining an accurate measurement. For example, adjustments may need to be made to the reading when the sample temperature deviates from the standard sample temperature, which is commonly 75-degrees-Fahrenheit. Green, immature beans will read a drier moisture than they actually are, so add 1.5%-points when you have these beans mixed in with sound beans.

When possible, harvest around the low-lying, frost-damaged grain areas, as this grain will have a high storage risk. Rather, harvest and handle them separately. Frost-damaged, low test-weight corn maintains most of its value for animal feed, but it probably won't be wanted by ethanol operations. Green soybeans are often discounted by processors.

Wet grain also has poor storage quality, so grain drying will be very important this year for those that plan to store through the winter. To safely store through the winter, dry corn to 15%-moisture and soybeans to 13%. Dry low test-weight corn and corn with damaged kernels to one percentage point lower in moisture content than normal.

High-temperature drying should be limited to 160-degrees for frost-damaged corn and to 130-degrees for soybeans to limit damage in the dryer. Use in-bin cooling rather than in-dryer to increase your dryer's capacity.

Low-temperature or natural-air grain drying should be limited to 21% moisture corn or dryer to prevent spoilage during drying. Natural-air drying is limited after late October; refer to equilibrium moisture content charts to determine if any drying will occur based on ambient air temperature and relative humidity. When average daily temperatures cool to below 40-degrees this fall, focus on getting the grain cooled for winter storage, aerate through the winter as needed, and continue drying when temperatures rise in the spring. Cool immature or damaged grain in bins to 30-40-degrees to store through the winter. Check grain for rising temperatures and moisture levels, odors, and insect activity every 2-4 weeks through the winter and every other week during spring.

It is especially important to clean and "core" bins with poor grain quality to remove fines accumulated in the center. Core the bin after filling it by removing about half of the peak height to improve aeration. Plan on checking lower quality stored grain frequently, aerating it as necessary, and market it before summer.

If we get wet conditions this fall that delay harvest, we may also want to look out for the development of molds and toxins, such as vomitoxin, on grain left out in the field.

### Grain Bin Safety:

When removing grain from a bin, visually inspect to be sure that an inverted cone has been created. If no cone is created, there may be bridging of the grain surface and a hollow space beneath. This is a very unsafe condition, so do not enter the bin until the bridging has been corrected. You should also not enter a bin when grain is flowing. Lastly, be sure to protect your lungs with proper respiratory protection when working with dusty or moldy grain.


**Figure:** U2U Corn GDD calculator for Johnson County, IA with 111-day corn for a planting date June 10.

#### Resources:

Frost Damage to Corn and Soybeans, PM-1635, Charles R. Hurburgh and Garren O. Benson, 2012.

High Moisture Corn Drying and Storage presentation, Kenneth Hellevang, 2014.

Soybean Drying and Storage, PM-1636, Charles R. Hurburgh, 2008.



## Thatch and Fall Removal

Article Author: Richard Jauron, Willy Klein

Thatch can be problematic and beneficial to lawns. Thatch supplies necessary food sources for microbes and organic matter. However, excessive thatch can harbor diseases and insects. Thatch that is properly balanced promotes growth and appearance of a well-groomed lawn. Horticulturists with ISU Extension and Outreach answer questions about thatch and fall removal. To have more questions answered, contact Hortline at [hortline@iastate.edu](mailto:hortline@iastate.edu) or 515-294-3108.



### What is thatch?

Thatch is the layer of dead and living plant material that accumulates above the soil surface in lawns. Thatch is composed primarily of shoots, crowns and roots. Contrary to popular belief, grass clippings do not contribute significantly to thatch buildup. When lawns are mowed properly, grass clippings break down quickly.

Thatch, to some degree, is present in all lawns. A small amount of thatch is beneficial as it moderates soil temperatures. However, thatch becomes detrimental when it is present in amounts greater than ½ inch. Excessive amounts of thatch increase the potential for turf damage due to drought, extremes in temperature, diseases and insects.

Compacted soils and heavy, clay soils are prone to thatch buildup. Heavy nitrogen fertilization and over-watering promote thatch accumulation.

### How do you remove thatch?

Thatch layers of ½ inch or greater can be removed with a vertical mower or power rake. The vertical mower has evenly spaced knife-like blades that rotate perpendicularly to the thatch surface. The vertical mower blades slice through the thatch and about ¼ inch into the soil. After vertical mowing, hand rake the loose thatch from the lawn. A powder rake has spring steel tines that loosen thatch without cutting into the soil. Vertical mowing and power raking services are available from professional lawn care companies. For individuals who prefer to do things themselves, vertical mowers and power rakes are often available at rental agencies.

### When is the best time to remove thatch from the lawn?

September is the best time to dethatch Kentucky bluegrass lawns. The slicing and ripping action during dethatching injures the grass. Dethatching the lawn in September gives the turfgrass several weeks of favorable weather in which to recover.

### How do I prevent the buildup of thatch?

Good lawn care practices can prevent the buildup of excessive amounts of thatch. Fertilize the lawn moderately. A single application of fertilizer in late October/early November is usually sufficient for low maintenance lawns. Highly maintained lawns may be fertilized in spring, September and late October/early November. Do not apply more than one pound of nitrogen per 1,000 square feet in a single application. If you decide to water the lawn in dry weather, apply 1 to 1½ inches of water per week in a single application or two applications three or four days apart. Core aerate lawns established on heavy, clay soils and soils that have become compacted.

---

## Are Those Around You Showing Signs of Stress?

Article Author: Tammy Jacobs, Human Sciences Extension and Outreach Hotlines Coordinator

Uncertainty in the farm economy makes this 24/7 resource extremely valuable

The [Iowa Concern Hotline](https://www.extension.iastate.edu/hotline) (800-447-1985) is a 24-hour a day, 7-day a week free, confidential resource for anyone with concerns or questions about farm finances, crisis and disaster response and personal health issues. Access to an attorney is also available to help provide legal education.



# Pesticide Applicator Training and Certification

Iowa State University Extension and Outreach, in cooperation with the Iowa Department of Agriculture and Land Stewardship (IDALS), provides training and certification programs for commercial and private pesticide applicators in Iowa. <https://www.extension.iastate.edu/psep/PSEP.html>

Certification procedures for pesticide applicators differ depending on the type of certification the applicator wishes to obtain.

All applicators are initially certified by passing an exam through IDALS.

## 2019 IDALS Pesticide Applicator Testing

Testing times at these locations are 10 a.m. – 2 p.m.

### Johnson County Extension Office

3109 Old Hwy 218 South Iowa City, IA 52246  
Phone: 319-337-2145

September 9, October 7, November 4, December 2

### Washington County Extension (Fairgrounds)

2223 – 250th St. Washington, IA 52353  
Phone: 319-653-4811

September 13, October 11, November 8, December 13

## Private Applicator

A person who applies any restricted-use pesticide for the production of an agricultural commodity on property owned or rented by the applicator or the applicator's employer, or on the property of other agricultural producers with whom the applicator trades services.

## Commercial Applicator

A person who applies pesticides or uses pest-control devices for hire on property other than his or her own.

To obtain initial certification as a commercial pesticide applicator, a person must pass both the closed-book exam over the [Iowa Core Manual IC 445](#), and an exam specific to the categories for which he or she needs to be certified to do the job.

See *IDALS Testing Schedule*

### Recertification by Exam

Recertify by passing exams, [covering the Iowa Core manual and appropriate category manuals] every 3 years. OR

### Recertify by attending a Commercial Continuing Instructional Course (C-CIC)

Fall Commercial Pesticide Applicator Continuing Instruction Courses			
Call your local county extension office to register for a course near you.			
Continuing Instruction Course	Date	Start	Cost
Roadside, Forest, and Aquatic Pest Management <i>Categories: 2, 5, 6, 10</i>	10/16/19	9 AM-11:30 AM	\$35 until 10/9, \$45 after 10/9
Mosquito and Public Health Pest Management <i>7D, 8, 10</i>	10/24/19	9 AM-11:30 AM	\$35 until 10/17, \$45 after 10/17
Ornamental and Turf Applicators <i>3O, 3T, 3OT, 10</i>	11/13/19	9 AM-11:30 AM	\$35 until 11/6, \$45 after 11/6
Fumigation <i>7C, 10</i>	11/19/19	9 AM-11:30 AM	\$35 until 11/12, \$45 after 11/12
Commercial Ag Weed, Insect, and Plant Disease Management <i>1A, 1B, 1C, 10</i>	11/20/19	9 AM-11:30 AM	\$35 until 11/13, \$45 after 11/13
Pest Control Operators <i>7A, 7B, 8, 10</i>	12/04/19	9 AM-11:30 AM	\$60 until 11/27, \$70 after 11/27
Greenhouse Tape <i>3G, 3O, 10</i>	Program Available October 1st.		\$35

If C-CICs are missed in one or more years of their three-year cycle, the applicator is required to take the certification exam(s) to renew his or her certification.

## Tri-State Ag Lender's Seminar

The 33rd Annual Tri-State Agriculture Lender's Seminar will be held on Thursday, October 24th at the Best Western Plus Hotel, 3100 Dodge Street, Dubuque, Iowa, from 9:30 am – 2:30 pm. This year's focus is on market outlooks for livestock, grains and dairy along with the issues of the farm technology, farm stress and tax laws affecting agriculture.

"Price risk management continues to be the major variable for profitability in many commodity enterprises. Understanding the ag markets is a necessary part of lender portfolio management", says Larry Tranel, ISU Extension Dairy Field Specialist." This seminar has proven itself in assisting lenders and farm financial advisors in helping farmers manage that risk."

The cost of the program is \$100 per person by October 14th. To register, contact Larry Tranel at ISU Extension in Dubuque County at 563-583-6496 or e-mail [tranel@iastate.edu](mailto:tranel@iastate.edu) for registration information.

## PROGRAM SPECIALISTS/ COORDINATORS Serving Region 15

### AGRICULTURE

#### Ag Engineering

Kristina Tebockhorst 319-337-2145  
ktebock@iastate.edu

#### Beef Livestock

Patrick Wall 641-842-2014  
patwall@iastate.edu

#### Commercial Horticulture

Patrick O'Malley 319-337-2145  
omall@iastate.edu

#### Agronomy

Rebecca Vittetoe 319-653-4811  
rka8@iastate.edu

#### Farm Management

Ryan Drollette 319-337-2145  
drollett@iastate.edu

#### Swine

Tom Miller 319-653-4811  
tmiller@iastate.edu

### COMMUNITY ECONOMIC DEVELOPMENT

Eric Christianson 515-231-6513  
ejchr@iastate.edu

### HUMAN SCIENCES

#### Family Life

Dawn Dunnegan 319-385-8126  
dunnegan@iastate.edu

#### Family Finance

Mary Weinand 641-472-4166  
mweinand@iastate.edu

#### Healthy Food Access Specialist

Jen Lamos 641-680-0052  
jenlamoss@iastate.edu

### 4-H YOUTH DEVELOPMENT

#### Youth Program Specialist-Field Team South

Sara Mohr 319-668-1052  
smohr@iastate.edu

### REGIONAL DIRECTOR

Gene Mohling 319-337-2145  
mohling@iastate.edu

### JOHNSON COUNTY STAFF

#### Office Coordinator

Brittnee Schneider 319-337-2145  
bschn@iastate.edu

#### 4-H County Youth Coordinator

Kate Yoder 319-337-2145  
yoder@iastate.edu

### Secure Pork Supply

In the event of an African Swine Fever (ASF), Foot and Mouth Disease (FMD), or Classical Swine Fever (CSF) outbreak in the United States, state and federal regulatory officials will restrict animal movement to slow or stop the spread of these foreign animal diseases (FADs). For livestock farms located in a disease control area, producers will need to meet certain criteria (for example: movement records and biosecurity measures in place) in order to request a permit to move animals. Guidance for requesting movement permits can be found in the Secure Pork Supply (SPS) Plan.

Iowa State University Swine Extension is helping producers understand what the state and federal response might include and how the resources in the SPS Plan can help prepare for a foreign animal disease outbreak. If we can help individual producers prepare, the industry as a whole is more prepared. Opportunities available include a variety of interactive workshops, webinars, and individual assistance on enhanced biosecurity plans and site maps.

Resources can be found on the SPS ([www.securepork.org](http://www.securepork.org)) and IPIC (<https://www.ipic.iastate.edu>) websites.

### New Youth Program Specialist Hired

Sara Mohr has been hired as Youth Program Specialist for Johnson and Iowa counties. Sara begins her position August 19. She will be housed in the Iowa County office.

Sara comes to Extension from the Benton Community Schools in Norway where she has been the At-Risk Coordinator/ Interventionist for 3 years and was a fourth grade teacher for 9 years. She has also been a Lead Induction Coach/Mentor and Professional Development Lead with the Grantwood Area Education Agency, Cedar Rapids. In these positions, she has developed, presented, and evaluated programming to fit all youth needs, advocated for under-represented youth, and mentored and coached educators. Sara received her MA from Coe College and BA from UNI. She has been a volunteer 4-H club leader.

Sara is very excited to work for ISU Extension and Outreach 4-H Youth Development where she hopes to have a great impact on the communities in Johnson and Iowa counties.

**IOWA STATE UNIVERSITY**  
Extension and Outreach

Johnson County Extension  
3109 Old Hwy 218 South  
Iowa City, IA 52246-5881

NONPROFIT ORG.  
U.S. POSTAGE  
PAID  
PERMIT #407  
IOWA CITY IA

RETURN SERVICE REQUESTED

**IOWA STATE UNIVERSITY**  
Extension and Outreach

**To continue  
receiving the Ag Newsletter in  
2020 please visit:**

<https://conta.cc/2lxAlxF>

Or contact Brittnee Schneider, Office Coordinator,  
at [bschn@iastate.edu](mailto:bschn@iastate.edu) or 319-337-2145