



Those with ACEs can be nearly twice as likely to be diagnosed with heart disease—the **#1 cause of death in America.**

EXPOSURE TO ACEs IS ALSO ASSOCIATED WITH INCREASED RISKS FOR:



STROKE:
2.1x as likely



DEPRESSION:
5.3x as likely



OBESITY:
1.2x as likely



CANCER:
1.4x as likely



CURRENT SMOKER:
3.1x as likely



ASTHMA:
2.2x as likely



KIDNEY DISEASE:
1.7x as likely



HEAVY DRINKER:
1.8x as likely



**HIGH SCHOOL
NON-COMPLETION**
1.4x as likely

CAMPAIGN FOR TRAUMA-INFORMED POLICY AND PRACTICE
WWW.CTIPP.ORG (WWW.CTIPP.ORG/NATIONALTRAUMACAMPAIGN)

Data based on: Merrick, M.T., Ford, D.C., Ports, K.A., Guinn, A.S., Chen, J., Klevens, J., Metzler, M., Jones, C.M., Simon, T.R., Daniel, V.M., Ottley, P., & Mercy, J.A. (2019). Vital signs: Estimated proportion of adult health problems attributable to adverse childhood experiences and implications for prevention — 25 states, 2015-2017. Centers for Disease Control and Prevention. Retrieved from <https://tinyurl.com/cdcaces19>